



Deep Reach Through Acid Encapsulation and Wormhole Propagation Sprint Project - DHRTC

Pandey, Jyoti Shanker; von Solms, Nicolas

Publication date:
2017

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Pandey, J. S., & von Solms, N. (2017). *Deep Reach Through Acid Encapsulation and Wormhole Propagation: Sprint Project - DHRTC*. Abstract from Danish Hydrocarbon Research and Technology Centre Technology Conference 2017, Lyngby, Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Danish Hydrocarbon Research and Technology Centre Technology Conference 2017

Deep Reach Through Acid Encapsulation and Wormhole Propagation Sprint Project - DHRTC

Jyoti Shanker Pandey (Research Assistant, CERE) (& Main Presenter) Prof Nicolas von Solms (Associate Professor-CERE)

Acid Injection into carbonate reservoirs is a popular well stimulation method to improve the formation permeability and enhance oil production. Acid injection leads to wormhole creation and propagation which can bypass the damage zone. Wormhole creation and propagation depends on the acid injection rate and dissolution rate of the rock in acid. At an optimal injection rate, dominant wormholes are formed which correspond to a minimum volume of acid required. Lower dissolution rates lead to deeper acid penetration into the formation.

The main objective of the project is to compare acid penetration and wormhole propagation in chalk samples (with and without cavity) using 3 different acidizing fluids (15% HCl, 10% Acetic Acid & Mixture of 15% HCl & 10% Acetic Acid). Initial test are carried out at normal temperature and pressure conditions. Key experiments performed are a soaking experiment, where acid is left stationary in a core and an acid injection experiment using a core flooding setup. Experiments are followed by CT Scanning, in order to elucidate the results.



AARHUS UNIVERSITY



UNIVERSITY OF
COPENHAGEN

Technical
University of
Denmark



GEUS



AALBORG UNIVERSITY
DENMARK